arXiv:1108.2409v1 [math.GR] 3 Aug 2011

Correction of a theorem on groups generated by transvections over $\mathbb{F}_2$

Hau-wen Huang

August 12, 2011

Abstract

Let V denote a finite dimensional vector space over the two-element field $\mathbb{F}_2$ endowed with a symplectic form B. Let $\operatorname{rad} V$ denote the radical of V with respect to B. Let $\mathrm{SL}(V)$ denote the special linear group of V. Let S denote a subset of V. Define $Tv(S)$ as the subgroup of $\mathrm{SL}(V)$ generated by the transvections with direction α for all $\alpha \in S$. Define $G(S)$ as the graph whose vertex set is S and where α, β in S are connected whenever $B(\alpha, \beta) = 1$. A well-known theorem states that if S spans V then $Tv(S)$ is isomorphic to a symmetric group if and only if $G(S)$ is a claw-free block graph. We give an example which shows that the necessity of the theorem is not true, and give a modification as follows. Let S denote a subset of $V \setminus \operatorname{rad} V$ and assume that S is a linearly independent set of V. If $Tv(S)$ is isomorphic to a symmetric group then $G(S)$ is a claw-free block graph.

1 A theorem on groups generated by transvections over $\mathbb{F}_2$

Throughout this note let V denote a finite dimensional vector space over the two-element field $\mathbb{F}_2$ endowed with a symplectic form B. Let $\operatorname{rad} V$ denote the radical of V with respect to B. Let $\mathrm{SL}(V)$ denote the special linear group of V. For $\alpha \in V$ define a linear transformation $\tau_\alpha : V \to V$ by

$$\tau_\alpha \beta = \beta + B(\beta, \alpha)\alpha \qquad \text{for all } \beta \in V.$$

We call τ_α the *transvection on V with direction α*. Observe that $\tau_\alpha^2 = 1$ and so $\tau_\alpha \in \mathrm{SL}(V)$. For a subset S of V, define $Tv(S)$ to be the subgroup of $\mathrm{SL}(V)$ generated by the transvections τ_α for $\alpha \in S$ and define $G(S)$ to be the simple graph which has vertex set S and an edge between vertices α and β if and only if $B(\alpha, \beta) = 1$.

A *claw* is a tree with one internal vertex and three leaves. A *claw-free graph* is a graph that does not have a claw as an induced subgraph. Let G denote a simple graph. A *cut-vertex of G* is a vertex whose deletion increases the number of components. A *block of G* is a maximal connected subgraph of G that has no cut-vertex. A *block graph* is a simple connected graph in which every block is a complete graph.

We now can state [3, Theorem 3.1] as follows.

Theorem 1.1. ([3, Theorem 3.1]). *Assume that S spans V. Then $Tv(S)$ is isomorphic to a symmetric group if and only if $G(S)$ is a claw-free block graph.*

2 A counterexample to the necessity of Theorem 1.1

In this section we show a counterexample to the necessity of Theorem 1.1. To this end we first recall some background material from [1, 2].

Definition 2.1. ([1, Section 3]). Define a binary relation $\mathcal{T}_0$ on the power set of V as follows. For any two subsets S and S' of V define $(S, S') \in \mathcal{T}_0$ whenever there exist $\alpha, \beta \in S$ such that S' is obtained from S by changing β to $\tau_\alpha \beta$.

Definition 2.2. ([1, Section 3]). Let $\mathcal{T}$ denote the equivalence relation on the power set of V generated by $\mathcal{T}_0$. The relation $\mathcal{T}$ is called the *t-equivalence relation.*

Lemma 2.3. ([2, Corollary 11.2]). *Let S denote a subset of $V \setminus \operatorname{rad} V$. Then S is t-equivalent to a subset S' of V whose associated graph $G(S')$ is a path if and only if $Tv(S)$ is isomorphic to a symmetric group.*

Example 2.4. Let V denote a vector space over $\mathbb{F}_2$ with dimension $n \geq 3$. Let $P = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ denote a basis for V. Define a symplectic form $B : V \times V \to \mathbb{F}_2$ by

$$B(\alpha_i, \alpha_j) = \begin{cases} 1 & \text{if } |i-j| = 1, \\ 0 & \text{if } |i-j| \neq 1. \end{cases}$$

Let $S = P \cup \{\alpha_1 + \alpha_2\}$. Note that S spans V. The set P can be obtained from S by changing $\alpha_1 + \alpha_2$ to $\tau_{\alpha_1}(\alpha_1 + \alpha_2) = \alpha_2$ hence $(S, P) \in \mathcal{T}$. Since $G(P)$ is a path and by Lemma 2.3, $Tv(S)$ is isomorphic to a symmetric group. The graph $G(S)$ is as follows.



Since the block of $G(S)$ with vertex set $\{\alpha_1, \alpha_2, \alpha_3, \alpha_1 + \alpha_2\}$ is not complete, $G(S)$ is not a block graph. Therefore we get a contradiction to the necessity of Theorem 1.1.

3 A modification of the necessity of Theorem 1.1

We modify the necessity of Theorem 1.1 as follows.

Theorem 3.1. *Let S denote a subset of $V \setminus \operatorname{rad} V$ and assume that S is a linearly independent set of V. If $Tv(S)$ is isomorphic to a symmetric group then $G(S)$ is a claw-free block graph.*

Rather than prove Theorem 3.1 by revising the original proof of the necessity of Theorem 1.1, we will provide a short proof. In our proof we will make use of the following definition and two lemmas.

Definition 3.2. Let $\mathcal{I}_0$ denote the restriction of $\mathcal{T}_0$ to the set of all linearly independent sets of V. Let $\mathcal{I}$ denote the equivalence relation on the set of all linearly independent sets of V generated by $\mathcal{I}_0$.

By checking the original proof of Lemma 2.3 carefully, we can get the following lemma.

Lemma 3.3. *Let S denote a subset of $V \setminus \operatorname{rad} V$ and assume that S is a linearly independent set of V. Then there exists a linearly independent set S' of V such that $(S', S) \in \mathcal{I}$ and $G(S')$ is a path if and only if $Tv(S)$ is isomorphic to a symmetric group.*

To state the following lemma we recall the notion of the line graph of a simple graph. Let G denote a simple graph. The *line graph $L(G)$ of G* is the simple graph that has a vertex for each edge of G, and two of these vertices are adjacent whenever the corresponding edges in G have a common vertex.

Lemma 3.4. ([4, Theorem 8.5]). *Let G denote a simple graph. Then G is a claw-free block graph if and only if G is the line graph of a tree.*

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. By Lemma 3.3 there exists a linearly independent set S' of V such that $(S', S) \in \mathcal{I}$ and $G(S')$ is a path. Observe that the binary relation $\mathcal{I}_0$ is symmetric. Therefore there exist some linearly independent sets $S_1 = S', S_2, \ldots, S_k = S$ of V such that $(S_1, S_2), (S_2, S_3), \ldots, (S_{k-1}, S_k) \in \mathcal{I}_0$. We now show that these graphs $G(S_i)$ for $1 \leq i \leq k$ are claw-free block graphs by induction on i. Note that $G(S_1)$ is a claw-free block graph. Let $2 \leq i \leq k$ be given. By induction hypothesis $G(S_{i-1})$ is a claw-free block graph. By Lemma 3.4 there exists a tree T_{i-1} that satisfies $L(T_{i-1}) = G(S_{i-1})$. Let $\alpha, \beta \in S_{i-1}$ such that the set S_i is obtained from S_{i-1} by changing β to $\tau_\alpha \beta$. If $B(\alpha, \beta) = 0$ then $S_{i-1} = S_i$ and there is nothing to prove. Thus we assume $B(\alpha, \beta) = 1$. Let u denote the common vertex of the edges α and β in T_{i-1}. Let v and w denote the other vertices adjacent to α and β in T_{i-1}, respectively. Let T_i denote the tree obtained from T_{i-1} by removing the edge β and adding a new edge between v and w. We call the new edge $\tau_\alpha \beta$. Observe that a vertex γ of $G(S_i)$ is adjacent to $\tau_\alpha \beta$ whenever γ is adjacent to exactly one of α and β in $G(S_{i-1})$. Therefore $L(T_i) = G(S_i)$ and so $G(S_i)$ is a claw-free block graph by Lemma 3.4. We have shown that $G(S_1), G(S_2), \ldots, G(S_k)$ are claw-free block graphs. The result follows. ☐

We finish this note with a comment. The original proof of the sufficiency of Theorem 1.1 does not use the assumption that S spans V. Actually we get the following result.

Theorem 3.5. *Let S denote a subset of V. If $G(S)$ is a claw-free block graph then $Tv(S)$ is isomorphic to a symmetric group.*

References

[1] R. Brown, S. P. Humphries. Orbits under symplectic transvections I. *Proceedings of the London Mathematical Society.* **52** (1986) 517–531.

[2] R. Brown, S. P. Humphries. Orbits under symplectic transvections II: the case $K = \mathbb{F}_2$. *Proceedings of the London Mathematical Society.* **52** (1986) 532–556.

[3] S. P. Humphries. Graphs and Nielsen transformations of symmetric, orthogonal and symplectic groups. *Quarterly Journal of Mathematics* (*Oxford*). **36** (1985) 297–313.

[4] F. Harary. *Graph Theory.* Addison-Wesley, Massachusetts, 1972.

Hau-wen Huang
Department of Applied Mathematics
National Chiao Tung University
1001 Ta Hsueh Road
Hsinchu, Taiwan, 30050.
Email: poker80.am94g@nctu.edu.tw